



Original Article

Quantitative Assessment of Chronotropic Incompetence Using Time Domain Heart Rate Variability Derived from 24 – Hours Ambulatory Holter Monitoring

Ardian Rizal^{1,2*}, Fitranti Suciati Laitupa³, Mohammad Saifur Rohman^{1,2}¹ Brawijaya Cardiovascular Research Center, Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Brawijaya, Malang, Indonesia.² Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Brawijaya, Malang, Indonesia.³ Department of Cardiology and Vascular Medicine, Dr. Iskak General Hospital, Tulungagung, Indonesia

ARTICLE INFO

Keywords:

Chronotropic incompetence;
Heart rate variability

ABSTRACT

Background : Chronotropic incompetence (CI) is defined as the failure of the heart to increase its heart rate along with activity and is an independent predictor of cardiovascular morbidity and mortality. The gold standard for CI is based on a treadmill test. However, up to now, there is no guideline to recommend whether CI could be concluded according to holter monitoring data.

Objective : This study aimed to determine CI based on Holter monitoring, specifically using time-domain analysis of heart rate variability (HRV).

Methods : This study used descriptive-analytic research with cross-sectional sampling. The population was patients that underwent treadmill and holter monitoring. The treadmill test was used as a gold standard of CI, and compare to holter monitoring using heart rate variability as a parameter. Data analysis used a comparative study (Independent T, Mann-Whitney, Chi-Square) and multivariate analysis logistic regression with a significance of $p < 0,05$.

Results: The subjects of the study were 111 patients with male-dominant (59 men). Standard Deviation N-N interval (SDNN) and Standard Deviation of Average N-N Interval (SDANN) were found lower in the CI group compared to the normal one. Mean SDNN was $113,57 \pm 51,08$ msec in CI group and $117,4 \pm 39,48$ msec in normal group with $p = 0.282$. Mean SDANN was $94,77 (73,42-118,85)$ in CI and $104,16 (74,9-139)$ with $p = 0.422$. While the Proportion of RR interval with the difference > 50 ms (RR50) and Average of All N-N Intervals (AVNN) found to be higher in CI group, but not statistically significant ($p = 0.681$; $p = 0.061$ respectively). Other parameter extracted from holter result, delta HR showed significant correlation with CI ($50.5 (43,25-61,5)$ vs. $56 (50-72)$ with $p = 0.014$).

Conclusion: Time-domain HRV didn't had a significant correlation with CI. However, delta HR from holter monitor could be used as a predictor of CI.

1. Introduction

Chronotropic Incompetence (CI) is often stated as the inability of the heart rate to increase in order to accommodate increasing cardiac demand.¹ More specifically, CI described as the inability to achieve 85% of maximum predicted heart rate adjusted to age, according to astrand formula.² Physiologically, the heart rate will increase when the body system told it to do so.³ A clear correlation between CI and increased mortality had been studied and established for decades.¹ Not only in patients with cardiovascular morbidity but also in those without any cardiovascular abnormality. Inability to reach 85%

of age-adjusted - predicted maximum heart rate is a predictor of mortality and future cardiovascular events, even after adjusting age and other cardiovascular risk factors (HR 1.75 %; CI 95%).⁴

The ability to reach the age-predicted maximal heart rate (APMHR) using a treadmill test is the gold standard diagnostic tool for CI. But it was traditionally formulated in healthy people. However, there was no standardized measurement in cardiac disease. In heart failure patients, the standard diagnostic formula is proven not accurate. Therefore, the modified formula has been proven superiorly accurate in heart failure patients who tacking beta-blocker although there was still

*Corresponding author at: Brawijaya Cardiovascular Research Center, Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Brawijaya, Malang, Indonesia
E-mail address: drardianrizal@ub.ac.id (A. Rizal).

<https://doi.org/10.21776/ub.hsj.2020.001.02.4>

Received 9 June 2020; Received in revised form 12 June 2020; Accepted 24 June 2020

Available online 18 July 2020

some limitation.⁵ Currently, many diagnostic approaches had been already proposed to make the diagnosis in certain cardiovascular conditions. But none of them showed a promising result.⁶

Holter monitoring had been widely used for diagnosing arrhythmia in clinical practice. Continuous monitoring of the patient can reflect the dynamic condition of the heart rate. Holter monitoring for diagnosing CI had been studied by lie et al in 2016. It showed that APMHR derived from a patient's daily activity could be used to diagnose CI.⁷ HRV is widely used to measure autonomic function. HRV is one parameter that could be extracted from holter result. In our clinical practice, we used to make non – objective or eyeballing evaluation of patient heart rate fluctuation during 24 hours to determine the chronotropic competency. In this study, we tried to analyze holter monitor results and made a quantitative measurement of chronotropic competence using holter monitor parameters, specifically HRV parameter.

2. Methods

2.1. Study Design

This was a single-center observational analytic study, with a cross-sectional design. The study took place in Saiful Anwar General Hospital, Cardiovascular Integrative Unit (Instalasi Pelayanan Jantung Terpadu / IPJT). The study was approved by Saiful Anwar General Hospital and Faculty of Medicine Universitas Brawijaya's ethical committee. Written informed consent was derived from all of the participants.

2.3. Study Participants

This study used purposive sampling for sample recruitment. We involved the patients that came to our cardiovascular rehabilitation unit who undergone a treadmill test procedure for various indications. They are asked for a willingness to did an additional 24 hours Holter examination. We exclude the patient with cardiac pacemaker and patient who had been previously diagnosed with the following condition: atrio-ventricular block; frequent premature ventricular contraction (PVC) and patient with atrial fibrillation (AF).

2.4. Data Collection and Measurements

A total of 117 patients was performing both treadmill test and Holter monitoring, but due to not meet the inclusion criteria, only 111 patients were complete by the end of the study.

All the patients underwent a treadmill test using whether bruce or modified bruce criteria. Chronotropic incompetence was defined as an inability to achieve 85 % age-predicted maximum heart rate (APMHR) or inability to achieve 0.6 of a chronotropic index in patients who take beta-blockers as a part of their medication. Chronotropic index (maximum HR during activity – resting HR) divided by (APMHR – resting HR) is a standardized measurement of increasing HR during exercise.⁸

Holter monitor was performed for 24 hours using 12 leads BTL cardio-point H 300. Standard holter measurements were recorded, such as maximum HR, minimum HR, delta HR, and any arrhythmia that being recorded. Additional time-domain HRV were automatically extracted from the data by the factory software. They are SDNN (Standard Deviation N-N interval), SDANN (Standard Deviation of Average N-N Interval), RMSDD (Root Mean Square of the Successive Differences), and pNNS50 (Number of Successive Differences between N-N interval that greater than 50 ms).

2.5. Statistical Analysis

Data were analyzed using SPSS for windows version 16. The comparison of CI and HRV parameters performed using paired student T-test if the data showed normal distribution or using Mann Whitney test if the data showed abnormal distribution.




Figure 1. Study flowchart

3. Results

3.1. Baseline Characteristics

From July 2017 until January 2018, there were approximately 900 treadmill procedures was done in the Cardiac Rehabilitation Centre of IPJT Saiful Anwar General Hospital. 117 patients willing to perform an additional 24 hours holter monitor procedure and joining the study. Six patients were excluded because of arrhythmia found in the holter result. Most of them were atrial fibrillation (5 patients). CI was found in 52 patients (46.9 %). Of those in CI group 50 % (26 patients) were male. While in normal group 55.93% (33 patients) were male. Diabetes mellitus were found more frequently in CI group (21.15 % vs 6.78 %, $p = 0.038$). Coronary artery disease, heart failure, smoker and beta-blocker were higher in CI group, although not statistically significant ($p = 0.49$; $p = 0.057$; $p = 0.45$ and $p = 0.13$ respectively).

3.2. Time domain HRV

From HRV parameters extracted automatically from holter result. None of all the parameters were significantly correlated with CI. SDNN and SDANN were found lower in CI group compared to the normal group. Mean SDNN was $113,57 \pm 51,08$ msec in CI group and $117,4 \pm 39,48$ msec in normal group with $p = 0.282$). Mean SDANN was $94,77$ (73,42-118,85) in CI and $104,16$ (74,9-139) with $p = 0.422$. While the RR50 and AVNN found to be higher in CI group, but not statistically significant ($p = 0.681$; $p = 0.061$ respectively).

The only holter parameter that found to be promising was delta HR. The difference between the maximum and minimum HR in CI group was 50.5 (43,25-61,5) versus 56 (50-72) with $p = 0.014$.

3.3. Multivariate Analysis

At the beginning of the study, we planned to make a model to quantitatively calculate the possibility of CI from time-domain HRV

Table 1. Bivariate Analysis of Categorical Variables

Variables	CI Group (n=52)	Normal Group (n=59)	p
Male	26 (50%)	33 (55,93%)	
Diabetes Mellitus	11 (21,15%)	4 (6,78%)	0.038
Heart Failure	19 (36,54%)	12 (20,34%)	0.057
Smoker	23 (44,23%)	22 (37,29%)	0.45
Using beta-blocker	30 (57,69%)	17 (28,81%)	0.13
Coronary Artery Disease	23 (44,23%)	31 (52,5%)	0.49

Note, data were presented in mean \pm SD or n(%)

Table 2. Bivariate Analysis of Numeric Variables

Variables	CI Group (n=52)	Normal Group (n=59)	p
Age	58,9 + 11, 46	55,7 + 11.2	0,138
SDNN (msec)	113,57 + 51,08	117, 4 + 39,48	0,282
SDANN (msec)	94,77 (73,42-118,85)	104,16 (74,9-139)	0,422
RR50 (msec)	5,51 (2,56-13,46)	4,95 (1,87-16,24)	0,681
AVNN (msec)	871,23 + 127,70	826,32 + 122,30	0,061
Delta HR (x/ menit)	50,5 (43,25-61,5)	56 (50-72)	0,014

Note, data were presented in mean \pm SD or n(%); SDNN, Standard Deviation N-N interval; SDANN, Standard Deviation of Average N-N Interval; RR50, Proportion of RR interval with the difference >50ms; AVNN, Average of All N-N Intervals ; HR, heart rate.

parameters, using logistic regression. The bivariate analysis p-value cut off was < 0.25. All parameters that had p-value < 0.25 will be eligible to join the next step analysis (logistic regression). Since there were only two holter parameters that had p-value < 0.25 (AVNN p = 0.061 and Delta HR p = 0.014), we decided not to continue to multivariate analysis.

4. Discussion

In this study, we found that only Delta HR is significantly correlated with CI (p 0.014). While other HRV parameters such as SDNN and SDANN were lower in CI group but not statistically significant. SDNN and SDANN reflect the sympathetic and parasympathetic activity. In patients with CI, HR failed to increase because of sympathetic activity was blunted. According to a previous study by Kawasaki et al, inability to reach APMHR in exercise (CI) is correlated with the ratio LF (Low Frequency) and HF (High Frequency), but not correlated with HF parameters.²⁰ Both HF and LF/HF ratios are the frequency domain HRV parameter. LF/HF ratio depicting sympathetic and parasympathetic, while HF is an index of parasympathetic tone.²¹ AVNN and Pnn50 found to be higher in CI group compared to non-CI, although not reached statistically significant. Unlike SDNN and SDANN, AVNN and pnn50 are more specific for a higher parasympathetic tone.

Delta HR, which is a simple measurement extracted from

holter results, also found to be a significant finding to predict CI. Patients with lower HR fluctuation found to be significantly correlated with the incidence of CI. Compared to eyeballing evaluation of HR fluctuation during 24 hours monitoring, delta HR offers more objective measurement. It opens the possibility for diagnosing CI using a currently popular and more simple wearable device, such as a smartwatch or smart band.

In this study, CI was found in 52 patients (46.8 %). According to ACC/AHA CI prevalence was ranged from 9 % to 89 %.¹ CI was found to be significantly higher in patients with diabetes mellitus type 2 (DMT2), heart failure, smoking, and bisoprolol therapy. Patients with DMT2 more likely to have CI, even after adjusting age, sex, and symptoms of CAD according to study among 12.291 patients.⁹ Diabetic autonomic neuropathy that common in DMT2 patients has been believed to be responsible for the development of CI in DMT2 patients. Hyperglycaemia will cause oxidative stress and inflammation through activation of several metabolic pathways, such as Protein kinase C (PKC), hexosamine, and increase of Advance glycation end products (AGE).¹⁰ It will end in systemic neuropathy, including autonomic neuropathy.¹¹

CI is a predictor of the increment of all-cause mortality in patients with DMT2. Patients with CI and DMT2 developed a higher risk (hazard ratio 2.3; with 95 % CI 1.8 -3.1) compare to those with DM only or CI only (hazard ratio 1.7).⁹

Prevalence of CI also higher in heart failure patients according to our study. This result also matched with another previous study. The prevalence was ranged between 25 -70 % in various studies.¹² Patients with heart failure are at a higher risk of developing CI, despite any concomitant beta-blocking agent therapy or not.¹³ The change in autonomic regulation, less contractility reserve, impaired physical activity, and beta-blocking agents contribute to the development of CI in HF patients.¹⁴

Data regarding the outcome of heart failure's patient with CI is not consistent. According to Yahya's study in 2010, CI was not a significant predictor of the worst outcome.¹⁵ While another study indicated that the inability to achieve APMHR was an independent predictor of mortality in heart failure.^{12,13,16}

Diagnosing CI in patients who take beta-blocker was a little bit different from those who don't. CI is common in patients with beta-blocker therapy.¹⁷ The result from a cohort study in 3.736 patients who take beta-blockers as routine medication, the presence of CI was an independent predictor of death the worst outcome.¹⁸ That's result came after adjusting age, gender, concomitant risk factor, and baseline heart rate. Discontinuation of the therapy didn't have a significant impact on reversing the CI in heart failure.¹⁹ In contrast, the weight loss program and exercise cardiac rehabilitation would improve CI in CAD patients who take beta-blockers.¹⁷

Study Limitation

We only used one out of three HRV measurements in this study. According to the 1996 Task Force of European Society of Cardiology (ESC) and the North American Society of Pacing and Electrophysiology (NAPSE), HRV can be measured using three methods, time-domain, frequency-domain, and geometric method⁽²²⁾. Only time-domain HRV was involved in the measurement. Further study using a combination of time-domain, frequency-domain, and geometric analysis hopefully will have more power in predicting CI.

5. Conclusion

Time-domain HRV can't be used to predict CI according to the final result. However, Delta HR which is a simple measurement that depicting heart rate fluctuation during the day had a strong relation to CI.

6. Declarations

6.1. Ethics Approval and Consent to participate

This study was approved by local Institutional Review Board, and all participants have provided written informed consent prior to involve in the study.

6.2. Consent for publication

Not applicable.

6.3. Availability of data and materials

Data used in our study were presented in the main text.

6.4. Competing interests

Not applicable.

6.5. Funding source

Not applicable.

6.6. Authors contributions

Idea/concept: AR. Design: AR. Control/supervision: MSR. Data collection/processing: AR, FSL. Extraction/Analysis/interpretation: AR, FSL. Literature review: MSR. Writing the article: AR. Critical review:

MSR. All authors have critically reviewed and approved the final draft and are responsible for the content and similarity index of the manuscript.

6.7. Acknowledgements

We thank to Brawijaya Cardiovascular Research Center.

References

1. Brubaker Peter H., Kitzman Dalane W. Chronotropic Incompetence. *Circulation*. 2011 Mar 8;123(9):1010–20.
2. Katritsis D, Camm AJ. Chronotropic incompetence: a proposal for definition and diagnosis. *Br Heart J*. 1993 Nov;70(5):400–2.
3. Astrand I. Aerobic work capacity in men and women with special reference to age. *Acta Physiol Scand Suppl*. 1960;49(169):1–92.
4. Lauer MS, Okin PM, Larson MG, Evans JC, Levy D. Impaired heart rate response to graded exercise. Prognostic implications of chronotropic incompetence in the Framingham Heart Study. *Circulation*. 1996 Apr 15;93(8):1520–6.
5. Efficacy and Safety of Exercise Training in Patients With Chronic Heart Failure: HF-ACTION Randomized Controlled Trial. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2916661/>
6. Zweerink Alwin, van der Lingen Anne-Lotte C.J., Handoko M. Louis, van Rossum Albert C., Allaart Cornelis P. Chronotropic Incompetence in Chronic Heart Failure. *Circ Heart Fail*. 2018 Aug 1;11(8):e004969.
7. Li H, Li J, Zhou X, Zhao J, Zhang F, Wu L, et al. Clinical feasibility of a new method to identify chronotropic incompetence in heart failure patients. *Physiol Meas*. 2016;37(11):1968–80.
8. Temporal Reduction in Chronotropic Index Predicts Risk of Cardiovascular Death Among Healthy Middle-Aged Men: a 28-Year Follow-Up Study | *Journal of the American Heart Association*. doi/10.1161/JAHA.116.004555
9. Azarbal B, Hayes S, Arbit B, Friedman J, Thomson L, Berman D. Relationship Between Diabetes Mellitus and Chronotropic Incompetence in Prediction of All-Cause Mortality. *J Am Coll Cardiol*. 2013 Mar 12;61(10 Supplement):E1115.
10. Verrotti A, Prezioso G, Scattoni R, Chiarelli F. Autonomic Neuropathy in Diabetes Mellitus. *Front Endocrinol*. 2014 Dec 1;5.
11. Balçioğlu AS, Müderrisoğlu H. Diabetes and cardiac autonomic neuropathy: Clinical manifestations, cardiovascular consequences, diagnosis and treatment. *World J Diabetes*. 2015 Feb 15;6(1):80–91.
12. Brubaker PH, Kitzman DW. Prevalence and management of chronotropic incompetence in heart failure. *Curr Cardiol Rep*. 2007 May;9(3):229–35.
13. Witte KKA, Cleland JGF, Clark AL. Chronic heart failure, chronotropic incompetence, and the effects of β blockade. *Heart*. 2006 Apr 1;92(4):481–6.
14. Zweerink A, van der Lingen A-L, Handoko M, Rossum A, Allaart C. Chronotropic Incompetence in Chronic Heart Failure: A State-of-the-Art Review. *Circ Heart Fail*. 2018 Aug 1;11.
15. Al-Najjar Y, Witte KK, Clark AL. Chronotropic incompetence and

- survival in chronic heart failure. *Int J Cardiol.* 2012 May 17;157(1):48–52.
16. Jorde UP, Vittorio TJ, Kasper ME, Arezzi E, Colombo PC, Goldsmith RL, et al. Chronotropic incompetence, beta-blockers, and functional capacity in advanced congestive heart failure: Time to pace? *Eur J Heart Fail.* 2008;10(1):96–101.
 17. Nugraha AW, Tjahjono CT, Rizal A. Increasing Age, Diabetes Mellitus and Beta Blocker Influence Heart Rate Recovery Values in Patients Undergoing Exercise Treadmill Test. *ACI Acta Cardiol Indones.* 2019 Mar 27;5(1):35–43.
 18. Khan MN, Pothier CE, Lauer MS. Chronotropic Incompetence as a Predictor of Death Among Patients With Normal Electrograms Taking Beta Blockers (Metoprolol or Atenolol). *Am J Cardiol.* 2005 Nov 1;96(9):1328–33.
 19. Hirsh B, Mignatti A, Garan A, Uriel N, Colombo P, Sims D, et al. Effect of -Blocker Cessation on Chronotropic Incompetence and Exercise Tolerance in Patients With Advanced Heart Failure. *Circ Heart Fail.* 2012 Aug 1;5:560–5.
 20. Kawasaki T, Kaimoto S, Sakatani T, Miki S, Kamitani T, Kuribayashi T, et al. Chronotropic incompetence and autonomic dysfunction in patients without structural heart disease. *Eur Eur Pacing Arrhythm Card Electrophysiol J Work Groups Card Pacing Arrhythm Card Cell Electrophysiol Eur Soc Cardiol.* 2010 Apr;12(4):561–6.
 21. Borchini R, Veronesi G, Bonzini M, Gianfagna F, Dashi O, Ferrario MM. Heart Rate Variability Frequency Domain Alterations among Healthy Nurses Exposed to Prolonged Work Stress. *Int J Environ Res Public Health.* 2018 Jan;15(1).
 22. Heart rate variability. *Eur Heart J.* 1996;17:28.